

10 Karmarkars Algorithm

We want to solve the following linear program:

- ▶ $\min v = c^t x$ subject to $Ax = 0$ and $x \in \Delta$.
- ▶ Here $\Delta = \{x \in \mathbb{R}^n \mid e^t x = 1, x \geq 0\}$ with $e^t = (1, \dots, 1)$ denotes the **standard simplex** in \mathbb{R}^n .

Further assumptions:

1. A is an $m \times n$ -matrix with rank m .
2. $Ae = 0$, i.e., the center of the simplex is feasible.
3. The optimum solution is 0 .

10 Karmarkars Algorithm

Suppose you start with $\max\{c^t x \mid Ax = b; x \geq 0\}$.

- ▶ Multiply c by -1 and do a minimization. \Rightarrow **minimization problem**
- ▶ We can check for feasibility by using the two phase algorithm. \Rightarrow **can assume that LP is feasible.**
- ▶ Compute the dual; pack primal and dual into one LP and minimize the duality gap. \Rightarrow **optimum is 0**
- ▶ Add a new variable pair x_ℓ, x'_ℓ (both restricted to be positive) and the constraint $\sum_i x_i = 1$. \Rightarrow **solution in simplex**
- ▶ Add $-(\sum_i x_i)b_i = -b_i$ to every constraint. \Rightarrow **vector b is 0**
- ▶ If A does not have full row rank we can delete constraints (or conclude that the LP is infeasible).
 \Rightarrow **A has full row rank**

We still need to make e/n feasible.

10 Karmarkars Algorithm

The algorithm computes **strictly feasible** interior points $x^{(0)} = \frac{e}{n}, x^{(1)}, x^{(2)}, \dots$ with

$$c^t x^{(k)} \leq 2^{-\Theta(L)} c^t x^{(0)}$$

For $k = \Theta(L)$. A point x is **strictly feasible** if $x > 0$.

If my objective value is close enough to 0 (the optimum!!) I can “snap” to an optimum vertex.

10 Karmarkars Algorithm

Iteration:

1. Distort the problem by mapping the simplex onto itself so that the **current point** \tilde{x} moves to the center.
2. Project the optimization direction c onto the feasible region. Determine a distance to travel along this direction such that you do not leave the simplex (and you do not touch the border). \hat{x}_{new} is the point you reached.
3. Do a backtransformation to transform \hat{x} into your new point \tilde{x}_{new} .

The Transformation

Let $\tilde{Y} = \text{diag}(\tilde{x})$ the diagonal matrix with entries \tilde{x} on the diagonal.

Define

$$F_{\tilde{x}} : x \mapsto \frac{\tilde{Y}^{-1}x}{e^{t\tilde{Y}^{-1}x}} .$$

The inverse function is

$$F_{\tilde{x}}^{-1} : \hat{x} \mapsto \frac{\tilde{Y}\hat{x}}{e^{t\tilde{Y}\hat{x}}} .$$

Note that $\tilde{x} > 0$ in every coordinate. Therefore the above is well defined.

Properties

$F_{\hat{x}}^{-1}$ really is the inverse of $F_{\hat{x}}$:

$$F_{\hat{x}}(F_{\hat{x}}^{-1}(\hat{x})) = \frac{\bar{Y}^{-1} \frac{\bar{Y}\hat{x}}{e^{t\bar{Y}\hat{x}}}}{e^{t\bar{Y}^{-1} \frac{\bar{Y}\hat{x}}{e^{t\bar{Y}\hat{x}}}}} = \frac{\hat{x}}{e^{t\hat{x}}} = \hat{x}$$

because $\hat{x} \in \Delta$.

Note that in particular every $\hat{x} \in \Delta$ has a preimage (Urbild) under $F_{\hat{x}}$.

Properties

\tilde{x} is mapped to e/n

$$F_{\tilde{x}}(\tilde{x}) = \frac{\tilde{Y}^{-1}\tilde{x}}{e^t\tilde{Y}^{-1}\tilde{x}} = \frac{e}{e^te} = \frac{e}{n}$$

A unit vectors e_i is mapped to itself:

$$F_{\bar{x}}(e_i) = \frac{\bar{Y}^{-1}e_i}{e^t \bar{Y}^{-1}e_i} = \frac{(0, \dots, 0, 1/\bar{x}_i, 0, \dots, 0)^t}{e^t (0, \dots, 0, 1/\bar{x}_i, 0, \dots, 0)^t} = e_i$$

All nodes of the simplex are mapped to the simplex:

$$F_{\bar{x}}(\mathbf{x}) = \frac{\bar{Y}^{-1}\mathbf{x}}{e^t \bar{Y}^{-1}\mathbf{x}} = \frac{\left(\frac{x_1}{\bar{x}_1}, \dots, \frac{x_n}{\bar{x}_n}\right)^t}{e^t \left(\frac{x_1}{\bar{x}_1}, \dots, \frac{x_n}{\bar{x}_n}\right)^t} = \frac{\left(\frac{x_1}{\bar{x}_1}, \dots, \frac{x_n}{\bar{x}_n}\right)^t}{\sum_i \frac{x_i}{\bar{x}_i}} \in \Delta$$

The Transformation

Easy to check:

- ▶ $F_{\bar{x}}^{-1}$ really is the inverse of $F_{\bar{x}}$.
- ▶ \bar{x} is mapped to e/n .
- ▶ A unit vectors e_i is mapped to itself.
- ▶ All nodes of the simplex are mapped to the simplex.

10 Karmarkars Algorithm

We have the problem

$$\begin{aligned} & \min\{c^t x \mid Ax = 0; x \in \Delta\} \\ &= \min\{c^t F_{\hat{x}}^{-1}(\hat{x}) \mid AF_{\hat{x}}^{-1}(\hat{x}) = 0; F_{\hat{x}}^{-1}(\hat{x}) \in \Delta\} \\ &= \min\{c^t F_{\hat{x}}^{-1}(\hat{x}) \mid AF_{\hat{x}}^{-1}(\hat{x}) = 0; \hat{x} \in \Delta\} \\ &= \min\left\{\frac{c^t \bar{Y} \hat{x}}{e^t \bar{Y} \hat{x}} \mid \frac{A \bar{Y} \hat{x}}{e^t \bar{Y} \hat{x}} = 0; \hat{x} \in \Delta\right\} \end{aligned}$$

Since the optimum solution is 0 this problem is the same as

$$\min\{\hat{c}^t \hat{x} \mid \hat{A} \hat{x} = 0, \hat{x} \in \Delta\}$$

with $\hat{c} = \bar{Y}^t c = \bar{Y} c$ and $\hat{A} = A \bar{Y}$.

Note that $e^t \bar{Y} x > 0$ for $x \in \Delta$.

We still need to make e/n feasible.

- ▶ We know that our LP is feasible. Let \bar{x} be a feasible point.
- ▶ Apply $F_{\bar{x}}$, and solve

$$\min\{\hat{c}^t x \mid \hat{A}x = 0; x \in \Delta\}$$

- ▶ The feasible point is moved to the center.

10 Karmarkars Algorithm

When computing \hat{x}_{new} we do not want to leave the simplex or touch its boundary (why?).

For this we compute the radius of a ball that completely lies in the simplex.

$$B\left(\frac{e}{n}, \rho\right) = \left\{x \in \mathbb{R}^n \mid \left\|x - \frac{e}{n}\right\| \leq \rho\right\} .$$

We are looking for the largest radius r such that

$$B\left(\frac{e}{n}, r\right) \cap \{x \mid e^t x = 1\} \subseteq \Delta.$$

10 Karmarkars Algorithm

This holds for $r = \left\| \frac{e}{n} - (e - e_1) \frac{1}{n-1} \right\|$. (r is the distance between the center e/n and the center of the $(n-1)$ -dimensional simplex obtained by intersecting a side ($x_i = 0$) of the unit cube with Δ .)

This gives $r = \frac{1}{\sqrt{n(n-1)}}$.


Now we consider the problem

$$\min\{\hat{c}^t x \mid \hat{A}x = 0, x \in B(e/n, r) \cap \Delta\}$$

This problem is easy to solve!!!

$$r^2 = (n-1) \cdot \left(\frac{1}{n} - \frac{1}{n-1} \right)^2 + \frac{1}{n^2} = \frac{1}{n^2(n-1)} + \frac{1}{n^2} = \frac{1}{n(n-1)}$$

The Simplex



10 Karmarkars Algorithm

Ideally we would like to go in direction of $-\hat{c}$ (starting from the center of the simplex).

However, doing this may violate constraints $\hat{A}\hat{x} = 0$ or the constraint $\hat{x} \in \Delta$.

Therefore we first project \hat{c} on the nullspace of

$$B = \begin{pmatrix} \hat{A} \\ e^t \end{pmatrix}$$

We use

$$P = I - B^t(BB^t)^{-1}B$$

Then

$$\hat{d} = P\hat{c}$$

is the required projection.

10 Karmarkars Algorithm

We get the new point

$$\hat{x}(\rho) = \frac{e}{n} - \rho \frac{\hat{d}}{\|\hat{d}\|}$$

for $\rho < r$.

Choose $\rho = \alpha r$ with $\alpha = 1/4$.

Iteration of Karmarkars Algorithm

- ▶ Current solution \bar{x} . $\bar{Y} := \text{diag}(\bar{x}_1, \dots, \bar{x}_n)$.
- ▶ Transform problem via $F_{\bar{x}}(x) = \frac{\bar{Y}^{-1}x}{e^t \bar{Y}^{-1}x}$.
Let $\hat{c} = \bar{Y}c$, and $\hat{A} = A\bar{Y}$.

- ▶ Compute

$$\hat{d} = (I - B^t(BB^t)^{-1}B)\hat{c} ,$$

where $B = \begin{pmatrix} \hat{A} \\ e^t \end{pmatrix}$.

- ▶ Set

$$\hat{x}_{\text{new}} = \frac{e}{n} - \rho \frac{\hat{d}}{\|\hat{d}\|} ,$$

with $\rho = \alpha r$ with $\alpha = 1/4$ and $r = 1/\sqrt{n(n-1)}$.

- ▶ Compute $\bar{x}_{\text{new}} = F_{\bar{x}}^{-1}(\hat{x}_{\text{new}})$.

Lemma 2

The new point \hat{x}_{new} in the transformed space is the point that minimizes the cost $\hat{c}^t \hat{x}$ among all feasible points in $B(\frac{\epsilon}{n}, \rho)$.

Proof: Let \hat{z} be another feasible point in $B(\frac{\epsilon}{n}, \rho)$.

As $\hat{A}\hat{z} = 0$, $\hat{A}\hat{x}_{\text{new}} = 0$, $e^t \hat{z} = 1$, $e^t \hat{x}_{\text{new}} = 1$ we have

$$B(\hat{x}_{\text{new}} - \hat{z}) = 0 .$$

Further,

$$\begin{aligned}(\hat{c} - \hat{d})^t &= (\hat{c} - P\hat{c})^t \\ &= (B^t(BB^t)^{-1}B\hat{c})^t \\ &= \hat{c}^t B^t(BB^t)^{-1}B\end{aligned}$$

Hence, we get

$$(\hat{c} - \hat{d})^t(\hat{x}_{\text{new}} - \hat{z}) = 0 \text{ or } \hat{c}^t(\hat{x}_{\text{new}} - \hat{z}) = \hat{d}^t(\hat{x}_{\text{new}} - \hat{z})$$

which means that the cost-difference between \hat{x}_{new} and \hat{z} is the same measured w.r.t. the cost-vector \hat{c} or the projected cost-vector \hat{d} .

But

$$\frac{\hat{d}^t}{\|\hat{d}\|} (\hat{x}_{\text{new}} - \hat{z}) = \frac{\hat{d}^t}{\|\hat{d}\|} \left(\frac{e}{n} - \rho \frac{\hat{d}}{\|\hat{d}\|} - \hat{z} \right) = \frac{\hat{d}^t}{\|\hat{d}\|} \left(\frac{e}{n} - \hat{z} \right) - \rho < 0$$

as $\frac{e}{n} - \hat{z}$ is a vector of length at most ρ .

This gives $\hat{d}(\hat{x}_{\text{new}} - \hat{z}) \leq 0$ and therefore $\hat{c}\hat{x}_{\text{new}} \leq \hat{c}\hat{z}$.

In order to measure the progress of the algorithm we introduce a **potential function** f :

$$f(\mathbf{x}) = \sum_j \ln\left(\frac{c^t \mathbf{x}}{x_j}\right) = n \ln(c^t \mathbf{x}) - \sum_j \ln(x_j) .$$

- ▶ The function f is invariant to scaling (i.e., $f(k\mathbf{x}) = f(\mathbf{x})$).
- ▶ The potential function essentially measures **cost** (note the term $n \ln(c^t \mathbf{x})$) but it penalizes us for choosing x_j values very small (by the term $-\sum_j \ln(x_j)$; note that $-\ln(x_j)$ is always positive).

For a point \hat{z} in the transformed space we use the potential function

$$\begin{aligned}\hat{f}(\hat{z}) &:= f(F_{\bar{x}}^{-1}(\hat{z})) = f\left(\frac{\bar{Y}\hat{z}}{e^t \bar{Y}\hat{z}}\right) = f(\bar{Y}\hat{z}) \\ &= \sum_j \ln\left(\frac{c^t \bar{Y}\hat{z}}{\bar{x}_j \hat{z}_j}\right) = \sum_j \ln\left(\frac{\hat{c}^t \hat{z}}{\hat{z}_j}\right) - \sum_j \ln \bar{x}_j\end{aligned}$$

Observation:

This means the potential of a point in the transformed space is simply the potential of its pre-image under F .

Note that if we are interested in **potential-change** we can ignore the additive term above. Then f and \hat{f} have the same form; only c is replaced by \hat{c} .

The basic idea is to show that one iteration of Karmarkar results in a constant decrease of \hat{f} . This means

$$\hat{f}(\hat{x}_{\text{new}}) \leq \hat{f}\left(\frac{e}{n}\right) - \delta ,$$

where δ is a constant.

This gives

$$f(\tilde{x}_{\text{new}}) \leq f(\tilde{x}) - \delta .$$


Lemma 3

There is a feasible point z (i.e., $\hat{A}z = 0$) in $B(\frac{e}{n}, \rho) \cap \Delta$ that has

$$\hat{f}(z) \leq \hat{f}\left(\frac{e}{n}\right) - \delta$$

with $\delta = \ln(1 + \alpha)$.

Note that this shows the existence of a good point within the ball. In general it will be difficult to find this point.



Let z^* be the feasible point in the transformed space where $\hat{c}^t x$ is minimized. (Note that in contrast \hat{x}_{new} is the point in the intersection of the feasible region and $B(\frac{e}{n}, \rho)$ that minimizes this function; in general $z^* \neq \hat{x}_{\text{new}}$)

z^* must lie at the boundary of the simplex. This means $z^* \notin B(\frac{e}{n}, \rho)$.

The point z we want to use lies farthest in the direction from $\frac{e}{n}$ to z^* , namely

$$z = (1 - \lambda) \frac{e}{n} + \lambda z^*$$

for some positive $\lambda < 1$.

Hence,

$$\hat{c}^t z = (1 - \lambda) \hat{c}^t \frac{e}{n} + \lambda \hat{c}^t z^*$$

The optimum cost (at z^*) is zero.

Therefore,

$$\frac{\hat{c}^t \frac{e}{n}}{\hat{c}^t z} = \frac{1}{1 - \lambda}$$

The improvement in the potential function is

$$\begin{aligned}\hat{f}\left(\frac{e}{n}\right) - \hat{f}(z) &= \sum_j \ln\left(\frac{\hat{c}^t \frac{e}{n}}{\frac{1}{n}}\right) - \sum_j \ln\left(\frac{\hat{c}^t z}{z_j}\right) \\ &= \sum_j \ln\left(\frac{\hat{c}^t \frac{e}{n}}{\hat{c}^t z} \cdot \frac{z_j}{\frac{1}{n}}\right) \\ &= \sum_j \ln\left(\frac{n}{1-\lambda} z_j\right) \\ &= \sum_j \ln\left(\frac{n}{1-\lambda} \left((1-\lambda)\frac{1}{n} + \lambda z_j^*\right)\right) \\ &= \sum_j \ln\left(1 + \frac{n\lambda}{1-\lambda} z_j^*\right)\end{aligned}$$

We can use the fact that for non-negative s_i

$$\sum_i \ln(1 + s_i) \geq \ln(1 + \sum_i s_i)$$

This gives

$$\begin{aligned} \hat{f}\left(\frac{e}{n}\right) - \hat{f}(z) &= \sum_j \ln\left(1 + \frac{n\lambda}{1-\lambda} z_j^*\right) \\ &\geq \ln\left(1 + \frac{n\lambda}{1-\lambda}\right) \end{aligned}$$

Suppose true for s_1, \dots, s_{k-1} . Then

$$\begin{aligned} \sum_{i=1}^k \ln(1 + s_i) &\geq \ln(1 + \sum_{i=1}^{k-1} s_i) + \ln(1 + s_k) = \ln\left((1 + \sum_{i=1}^{k-1} s_i)(1 + s_k)\right) \\ &= \ln\left(1 + \sum_i s_i + s_k \sum_{i=1}^{k-1} s_i\right) \geq \ln(1 + \sum_i s_i) \end{aligned}$$

In order to get further we need a bound on λ :

$$\alpha r = \rho = \|z - e/n\| = \|\lambda(z^* - e/n)\| \leq \lambda R$$

Here R is the radius of the ball around $\frac{e}{n}$ that contains the whole simplex.

$R = \sqrt{(n-1)/n}$. Since $r = 1/\sqrt{(n-1)n}$ we have $R/r = n-1$ and

$$\lambda \geq \alpha \frac{r}{R} \geq \alpha/(n-1)$$

Then

$$1 + n \frac{\lambda}{1-\lambda} \geq 1 + \frac{n\alpha}{n-\alpha-1} \geq 1 + \alpha$$

This gives the lemma.

Lemma 4

If we choose $\alpha = 1/4$ and $n \geq 4$ in Karmarkars algorithm the point \hat{x}_{new} satisfies

$$\hat{f}(\hat{x}_{\text{new}}) \leq \hat{f}\left(\frac{e}{n}\right) - \delta$$

with $\delta = 1/10$.

Proof:

Define

$$\begin{aligned}g(\hat{x}) &= n \ln \frac{\hat{c}^t \hat{x}}{\hat{c}^t \frac{e}{n}} \\ &= n(\ln \hat{c}^t \hat{x} - \ln \hat{c}^t \frac{e}{n}) .\end{aligned}$$

This is the change in the **cost part** of the potential function when going from the center $\frac{e}{n}$ to the point \hat{x} in the **transformed space**.

Similar, the **penalty** when going from $\frac{e}{n}$ to w increases by

$$h(\hat{x}) = \text{pen}(\hat{x}) - \text{pen}\left(\frac{e}{n}\right) = - \sum_j \ln \frac{\hat{x}_j}{\frac{1}{n}}$$

where $\text{pen}(v) = - \sum_j \ln(v_j)$.

We want to derive a lower bound on

$$\begin{aligned}\hat{f}\left(\frac{e}{n}\right) - \hat{f}(\hat{x}_{\text{new}}) &= [\hat{f}\left(\frac{e}{n}\right) - \hat{f}(z)] \\ &\quad + h(z) \\ &\quad - h(\hat{x}_{\text{new}}) \\ &\quad + [g(z) - g(\hat{x}_{\text{new}})]\end{aligned}$$

where z is the point in the ball where \hat{f} achieves its minimum.

We have

$$[\hat{f}(\frac{e}{n}) - \hat{f}(z)] \geq \ln(1 + \alpha)$$

by the previous lemma.

We have

$$[g(z) - g(\hat{x}_{\text{new}})] \geq 0$$

since \hat{x}_{new} is the point with minimum cost in the ball, and g is monotonically increasing with cost.

We show that the change $h(w)$ in **penalty** when going from e/n to w fulfills

$$|h(w)| \leq \frac{\beta^2}{2(1-\beta)}$$

where $\beta = n\alpha r$ and w is some point in the ball $B(\frac{e}{n}, \alpha r)$.

Hence,

$$\hat{f}\left(\frac{e}{n}\right) - \hat{f}(\hat{x}_{\text{new}}) \geq \ln(1 + \alpha) - \frac{\beta^2}{(1 - \beta)} .$$

Lemma 5

For $|x| \leq \beta < 1$

$$|\ln(1+x) - x| \leq \frac{x^2}{2(1-\beta)} .$$

For $|x| < 1$

$$\ln(1+x) = \sum_{i \geq 1} (-1)^{i+1} \frac{x^i}{i} = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$$

This gives

$$\begin{aligned} |\ln(1+x) - x| &\leq \left| -\frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots \right| \leq \left| \frac{x^2}{2} + \frac{x^3}{3} + \frac{x^4}{4} + \dots \right| \\ &\leq \frac{x^2}{2} |x^0 + x^1 + x^2 + \dots| = \frac{x^2}{2(1-|x|)} . \end{aligned}$$

This gives for $w \in B(\frac{\epsilon}{n}, \rho)$

$$\begin{aligned} |h(w)| &= \left| \sum_j \ln \frac{w_j}{1/n} \right| \\ &= \left| \sum_j \ln \left(\frac{1/n + (w_j - 1/n)}{1/n} \right) - \sum_j n \left(w_j - \frac{1}{n} \right) \right| \quad [= 0] \\ &= \left| \sum_j \left[\ln \left(1 + \underbrace{n(w_j - 1/n)}_{\leq n|x| < 1} \right) - \underbrace{n(w_j - 1/n)}_x \right] \right| \\ &\leq \sum_j \frac{n^2(w_j - 1/n)^2}{2(1 - \alpha nr)} \\ &\leq \frac{(\alpha nr)^2}{2(1 - \alpha nr)} \end{aligned}$$

The decrease in potential is therefore at least

$$\ln(1 + \alpha) - \frac{\beta^2}{1 - \beta}$$

with $\beta = n\alpha r = \alpha\sqrt{\frac{n}{n-1}}$.

It can be shown that this is at least $\frac{1}{10}$ for $n \geq 4$ and $\alpha = 1/4$.

Let $\tilde{x}^{(k)}$ be the current point after the k -th iteration, and let $\tilde{x}^{(0)} = \frac{e}{n}$.

Then $f(\tilde{x}^{(k)}) \leq f(e/n) - k/10$.

This gives

$$\begin{aligned} n \ln \frac{c^t \tilde{x}^{(k)}}{c^t \frac{e}{n}} &\leq \sum_j \ln \tilde{x}_j^{(k)} - \sum_j \ln \frac{1}{n} - k/10 \\ &\leq n \ln n - k/10 \end{aligned}$$

Choosing $k = 10n(\ell + \ln n)$ with $\ell = \Theta(L)$ we get

$$\frac{c^t \tilde{x}^{(k)}}{c^t \frac{e}{n}} \leq e^{-\ell} \leq 2^{-\ell} .$$

Hence, $\Theta(nL)$ iterations are sufficient. One iteration can be performed in time $\mathcal{O}(n^3)$.